

Qkine GDNF DISCs - Standard

Product Information Sheet




StemCultures | +1 518 621 0848 | support@stemcultures.com | www.stemcultures.com | www.qkine.com

Product Description

Qkine GDNF DISCs™ are patented inert, non-degradable, biocompatible hydrogels that release defined levels of GDNF stably into culture medium over the course of two weeks. Qkine GDNF DISCs™ are loaded with StemBeads® Qkine GDNF (Qk-SBGD1), PLGA microbeads that encapsulate GDNF (Qk051). GDNF is key for supporting neuronal differentiation. Qkine GDNF DISCs™ are easy to add and remove, giving scientists enhanced control of growth factor levels in their cultures. Controlled delivery and stable levels overcome the 0.5 hour half-life (Figure 1) of GDNF and improve cell cultures while saving researchers valuable time and resources.

Qkine GDNF DISCs™ have been tested in multiple neuronal base media, including Brain Phys, with enhanced cellular profiles. Qkine GDNF DISCs™ can be used with other StemCultures products.

Figure 1



Product Information

Catalog #	Product Name	Storage	Expiration	DISC™ Size	Recommended Well/Plate Size
Qk-DSCGD1-12	Qkine GDNF DISC™ (Standard)	4°C	4 months from manufacture (see label)	2-3 mm diameter, dry 5-6 mm diameter, rehydrated	6 well, 12 well

Directions for Use


NOTE: For 2D cultures, place DISC™ in a Transwell® or coat your plate with PDL and matrigel to avoid cell lifting. For 3D cultures (i.e. organoids) place DISC™ directly in the culture well.

- Using aseptic cell culture technique, wipe DISC™ container with 70% ethanol and place into a biosafety cabinet before opening.
- Plate cells and add culture medium to wells according to your typical protocol. Coat your plate if desired.
- Using sterile forceps, transfer each DISC™ into a culture well or Transwell® containing 2 mL of medium (see Release Data section on page 2 for more information on release and adjustments).

Note: As DISCs™ rehydrate, they will swell and become transparent. Embedded StemBeads® will be visible under a microscope. An image of a DISC™ under a microscope is on page 2 for reference.

- Replace the medium based on your typical feeding schedule. You can use a low powered vacuum or a pipette to remove the medium but not the DISC™. The original DISC™ can remain in your cultures for 2 weeks.
- When removing the DISC, we recommend removing the DISC before removing the medium.

Recommended Culture Schedule



Note: Different celllines, culture densities, and media may require adjusted schedules.

Please reach out to support@stemcultures.com for ordering and technical support.

Release Data

We recommend a release of 5 ng/mL when culturing neuronal cells. However, to fit other needs, the release can be adjusted slightly based on the amount of medium and the number of DISCs™ that are added. See the chart below for reference.

DISC™ Size	Volume of Medium Added	Number of DISCs™ per well	Release in Volume of Medium Added	Example Plate Size
Standard	2 mL	1	5 ng/mL	6 well
Standard	1 mL	1	10 ng/mL	12 or 24 well
Standard	0.5 mL	1	20 ng/mL	24 well
Standard	2 mL	2	10 ng/mL	6 well
Standard	2 mL	3	15 ng/mL	6 well
Standard	2 mL	4	20 ng/mL	6 well